A TWO-YEAR STUDY OF BIRDS COLONIZING RESTORED GRASSLANDS ON THE EASTERN SHORE OF MARYLAND - WITH SPECIAL FOCUS ON THE GRASSHOPPER SPARROW (*Ammodramus savannarum*)

27 April 2001

Gary J. Dodge[1], Brian Byrnes[2], and Douglas E. Gill[1]

[1]Program for Sustainable Development and Conservation Biology
Department of Biology, University of Maryland, College Park, MD 20742
Chester River Field Research Center, Chestertown MD
[2]Department of Biology, Swarthmore College, Swarthmore PA 19104

6
The Conservation Reserve Program (CRP) and Conservation Reserve Enhancement Program (CREP) ..6
Block and Strip Plantings 6
CRFRC at Chino Farms 7
III. 9
Nest Ecology and Studies..... *10*
Reproductive Success..... *11*
Territory Studies *12*

II. INTRODUCTION

regionally

Grassland Restoration on Maryland's Eastern Shore

The Conservation Reserve Program (CRP) and Conservation Reserve Enhancement Program (CREP)

The Conservation Reserve Program (CRP) of the USDA was enacted as part of the 1985 Farm Bill and was established to remove marginal agriculture land from production to reduce soil erosion and chemical runoff into wetlands and aquatic ecosystems and to improve wildlife habitat. The program provides monetary incentives for farmers to participate by converting some of their lands to trees, grasses, or other stabilizing vegetation. CRP pays an annual rental for the land as well as provides funds for conversion (i.e. seed, seedlings, special planting equipment, etc.). Contracts are written for 10 or 15 years and administered through state and local government agencies. CRP has put a high priority on the restoration of warm-season grasslands.

The Maryland Conservation Reserve Enhancement Program (CREP) is a partnership between Maryland and the federal government, and is federally sponsored through the CRP. In 1997 Maryland launched CREP with provision of $195 million for conversion of marginal agriculture lands on or adjacent to wetlands or waterways to natural wildlife habitats. The program received a significant boost with Vice President Al Gore's 1997 announcement of CREP's mandate to protect and stabilize up to 40,469 hectares (100,000 acres oflan0(i)-2 oflenzingheCesaheahed

relief of the Eastern Shore and farmland, very little suitable habitat on the Eastern Shore will meet this qualification.

Although CRP and CREP were designed with a major goal of establishing wildlife habitat, analyses of the effect of block CRP grasslands on grassland birds in the Midwest have yielded mixed results. Some CRP fields are reported to have positive effects on grassland birds in the Midwest (Best *et al*. 1998, Herkert 1998), but other studies reported no effect, mixed effects, or negative growth in other CRP fields (Reynolds *et al*. 1994, McCoy *et al*. 1999). There is little scientific evidence that the filter/buffer strips established by both CRP and CREP have had any effect on Grasshopper Sparrows or most other grassland birds (Delisle *et al*. 1996 as reported in Vickery 1996, Herkert 1998). Grasshopper Sparrows have been shown to favor large grassland areas and do not responded to linear habitats (Johnson and Temple 1986a, Warner 1992, Herkert 1994a, Vickery *et al.* 1994, Smith 1997). Herkert (1994) suggested that 12.1 hectares (30 acres) is the minimum area needed to sustain a breeding population in Illinois, but Grasshopper Sparrows were more likely to be established in large fragments of 130 – 146 hectares (320 – 360 acres) in Minnesota (Johnson and Temple 1986a), and colonization reached 50% incidence of 90 grassland sites in Maine at about 100 ha (Vickery *et al*. 1994). Although CREP strip acreage may provide cover for some birds, it also may provide cover for ed

In March 1999, the 93-hectare (230-acres) block was planted with mixtures of nine species of native perennial bunch grasses (Table 3). The 93 hectares were divided into twelve fields of approximately 8.1 hectares (20 acres) each and separated by firebreaks planted in clover. Each field was planted in a mixture of three species of grass. Five treatments (mixtures) with two replicates were created in Fields 1-10 (Fields 11 and 12 were not included in the experimental design). The planting design is shown in Table 4. Called the Experimental Grassland Restoration Project (on-site roadside signs refer to it as Grassland Plantation), this is the largest block of contiguous warm-season grasslands presently on the Eastern Shore of Maryland.

III. GRASSHOPPER SPARROW STUDIES

Introduction

We conducted comprehensive ecological studies on the Experimental Grassland Restoration Project of CRFRC during May through August of both summer 1999 and 2000. Due to resource limitations and the apparent concentration of Grasshopper Sparrow territories in Fields 1-8, we limited our study to these contiguous restored grassland blocks and only occasionally explored Fields 9-12. Summer 2001 will be the third year of the critical establishment period for these grasslands. Our 1999 and 2000 studies focused on –

1) the population responses of bird species of special concern, including Grasshopper Sparrow, Dickcissel, Bobolink, and Bobwhite Quail, to the restored grasslands, with particular attention to a comparison of the breeding performance of Grasshopper Sparrows in the five different treatments of plantings,
2) the growth performance of the eight species of native Warm Season Grasses planted in March 1999 and the complex plant community of alien and native species already present

(T) were placed above (proximal) to the USFWS band on the bird's left leg to designate males; Hot Pink (P) or Light Pink (F) was placed above the USFWS band on the left leg of females. Juveniles caught in mist nets were banded with only a federal aluminum band on the left leg. Nestlings from a common nest (siblings) were marked with a common colored band on their right leg to denote their nestship and an aluminum band on their right leg.

We used Student's T-Tests to compare morphometric differences between males and females.

Nest Ecology and Studies

Most Grasshopper Sparrow nests were discovered incidentally while conducting censuses, operating mist nets, and mapping territories in the fields. We occasionally flushed adult birds from nests during the course of our research activities, usually when we approached within five meters of a nest. Flushed adults often behaved with a broken-wing display as the bird ran and vocalized chip notes. In most cases the adults would persistently chip while researchers were in the immediate nest area. Early in 1999 we systematically searched the fields by walking in pairs 4-5 meters apart, but this method proved unproductive. On occasion we found nests by following adult birds carrying food to the nest until they dropped from a perch; we then rushed into the area to force a flush from the nest. Nevertheless, the nests were difficult to find and often required several minutes to find following the flush.

Once a flush was noted, the spot was marked with a hat or flag, and we would systematically search the area. After the nest was located, a strip of flagging was tied to a prominent piece of vegetation no closer than seven meters to the nest (most often 10 meters) and a nest record card was created with directions and a drawing to the nest from the flagging. We used a Garmin 12XL GPS receiver to note locations of both the flag and the nest.

We visited nests every other day to monitor the progress of development but not often enough to interfere with the parenting or to attract predators. Our routine was to approach the flags that marked nests from opposite direction of the nest and then visually locate the nest site from the flag before approaching. Nests were then checked for adult presence, number of eggs, chicks, and age of the chicks. If chicks or eggs were present (and weren't ready to be banded), researchers would leave the area quickly to minimize disturbance. If the nest was empty, it was assessed for signs of predation or fledging.

Nests were considered successful if at least one chick fledged. Fledging was determined by age and development of the chicks at the previous observation, and condition of the nest. Fecal sacs in and around the nest site were further evidence in support of fledging.

Whether a population was a source of recruits for the region or a local reproductive sink was

IV.

Grasshopper Sparrow Reproductive Studies

Nest Site Choice

In 2000 Grasshopper Sparrows chose nest sites that averaged 34% bare ground, 36% duff, and 30% live vegetation in a one-meter radius circle around the nest (Table 7). Bare ground and duff ranged from 5% to 70% cover, and live vegetation ranged from 15% to 65% in the area five meters around the nest.

The literature reports that Grasshopper Sparrows favor a moderate amount of bare ground (Whitmore 1979, Bollinger 1995). We have evidence that nest sites were chosen with a bias toward sparse vegetation. In Field 7 we found two nests, one in [illegible] and one in [illegible]

Our priorities at the onset of both the 1999 and 2000 seasons were to get the team of researchers trained and the adult population of Grasshopper Sparrows banded. We therefore did not investigate return dates and the establishment of territories as thoroughly as wished. As noted

to a daily probability of nest survival of .973 and a probability of nest survival to fledging of 56.3%.

Estimates of fecundity are shown in Table 11. In 1999, the average number of eggs laid per nest was 3.8 (n=5 nests) and average number of chicks per successful nest was 3.8 (n=7 nests). In 2000, the average number of eggs laid per nest was 4.8 (n=21 nests) and average number of chicks per successful nest was 4.4 (n=22 nests). Pooled over both years the average number of eggs per nest was 4.6 (n=27 nests) and the average number of chicks per successful nest was 4.3 (n=29 nests). These estimates of nest production are consistent with those reported by Vickery (1996). We haven't yet determined how many offspring each pair fledges in the breeding season and whether there are significant differences among adult age groups.

Dispersal of Hatching-year Grasshopper Sparrows

A total of twelve Local (banded in the nest) and six HY (hatching year – post fledge birds in juvenal plumage) Grasshopper Sparrows were banded in 1999 but none were ever observed again. In 2000 84 Local and 46 HY birds were banded, but none of these HY birds were seen again in 2000. We hope that some of these 130 birds will be found in 2001 and give us insight into the survival, dispersal, and behavior of the first-year/nestling birds from the CRFRC grasslands. The lack of repeat observations for the total 148 HY birds has several possible explanations: 1) fledglings instantly dispersed from their natal patch and established home bases (pre-SY territories) in new but distant habitat patches (J. Sheppard, pers. comm.); 2) all fledglings fell prey to predators – the Grassland Restoration fields could be a reproductive sink; 3) fledgling behavior was such that they eluded mist nets and observation.

Despite our failure to relocate any of our marked HY birds, we observed and captured numerous unbanded HY birds in the project fields in August of both years. The origin of these unbanded HY birds is intriguing. Clearly, they were either fledged from undiscovered nests in our grasslands or they were immigrants from other localities. The abundance of the unmarked HY

territory, recapture data indicated these females were traveling over multiple territories in short periods of time. The function of these travels is unknown (e.g. whether extra-pair copulations were achieved). We have no evidence yet that these females were actively nesting at the time, but it was during the height of the breeding season. We must note that our sample is biased to individuals that move more - more active females will be more likely caught in mist nets.

extraordinarily high pitches compared to other
xclusively in the 8 kHz range (Rising 1996, Vickery
some other sparrows, auditory sensitivity is at a
y above 6 kHz (Dooling *et al*. 2000, Fay 1988,
y in audibility with distance is greater (steeper)
encies. This raises the question: How do
e singing male itself hear advertising songs? At
es of Grasshopper Sparrows. Because bird hearing is
. 2000), we presume that auditory sensitivity of
small birds. Thus, the extra high frequency of the
on the distances over which they can communicate.
ds have two choices, either the power (decibels) of
ng carries a normal distance, or Grasshopper
e closer to neighbors. The fact that our preliminary
grassland birds supports the latter hypothesis.

tion in grasshopper sparrows in order to
es. Specifically, we will:

arrows with respect to the measurable physical
audiotape recordings to examine the relationship
ween-individual variation for birds sampled from

ing sounds in the restored grasslands. Examine
acteristics of these sounds to those of Grasshopper

VI. MANAG EMENT OPTIONS AND RECOMMENDATIONS

Bryan and Best 1991, Warner 1992, Kershner and Bollinger 1996, Vickery 1996). Grasslands on the East Coast may be more productive than in the Midwest (Bollinger 1995, Mitchell *et al*. 2000) and mowing without haying may increase litter. As mentioned previously, deeper litter can be a negative feature of Grasshopper Sparrow habitat (Bollinger 1995). Recommended times for mowing would be in August, after the majority of the nesting season, and haying would be advisable.

In the first two years of establishment of the Experimental Grassland Restoration Project, mowing has been a necessary technique to reduce weed competition for establishment of warm-season grasses. CRP and CREP administrators require this short-term management for enrolled acreage. Once the perennial grasses have become established, mowing is not recommended as a management tool for native prairie restoration due to increased litter buildup.

Increasing Habitat

Grasshopper Sparrows are loosely colonial and small populations appear to be somewhat ephemeral (Smith 1968, Holmes 1996). Population Viability Analyses show that populations of Grasshopper Sparrows show clear relationships between carrying capacity and population persistence (Wells 1997; Dodge, unpublished). Larger habitat carrying capacities would allow populations to grow large, better withstand variability in reproductive success, and buffer against environmental stochasticity. Increasing habitat available to Grasshopper Sparrows in the form of large habitat blocks, like those at CRFRC, would certainly increase the chances of the population persistence.

The apparent vigor with which Grasshopper Sparrows colonize new habitat blocks provides strong support for increasing habitat as a crucial management component. At our study site, two months after the conversion of marginal croplands to grasslands the population of Grasshopper Sparrows in the vicinity increased from approximately 10 mating pairs (J. Gruber, pers. comm.) to 75 mating pairs. We surmise that some of the 65 new males in the area were returning from the past breeding season, but that most were new immigrants. The implication is that site fidelity is not necessarily as strong in males as others (Welarreh(di)2(n)-8(g)12(sv -1.15 Td ( )T[])-2(e)4(c)-6(e)4(s)-1(s

VII. DICKCISSELS AT CRFRC

Results and Discussion

covers in Kansas (Hughes *et al*. 1999) but a positive association was found in Iowa (Bryan and Best 1991). Dickcissel reproductive success was positively associated with litter depth in Kansas CRP fields (Hughes *et al*. 1999).

Management of the GrasslaP 0 Td (. 199)Tc -0a.60 Td tor(e). 19C /CP thojntlds (H- Tc 0.004 Tw [(et)-6(1

VIII. OTHER BIRDS AND THEIR USES OF THE CRFRC GRASSLANDS

Killdeer were seen regularly early in both 1999 and 2000. A single nest was found in 1999.

Upland Sandpipers (a flock of 13) were observed in the study fields in August 1999. We assumed this to be a migrant group. None were observed in 2000.

Northern Bobwhite (three coveys and many adults) were observed in the fields in 1999 and 2000. A single observed nest was destroyed by a mower. Three coveys were present in late September 2000.

Eastern Kingbirds were abundant in the fields throughout both seasons. Most likely they are nesting on field borders.

Bobolinks (eight males and two females) were observed foraging in fields on 15 July 2000. Smaller groups were observed in the fields for about a week following.

Red-winged Blackbirds are abundant in fields for the entire season in both 1999 and 2000. Two nests were discovered in the fields in 2000.

Eastern Meadowlarks (a single pair) were regularly observed on Kibler Road about 1 mile from the restored grasslands in 1999. One male was singing in the fields for about a week in early June 2000.

Orchard Orioles were commonly observed foraging in fields in both 1999 and 2000. They are common breeders in the surrounding trees and in the shrub islands in the fields.

Northern (Baltimore) Orioles are common breeders in the surrounding trees and occasionally observed foraging in the fields in both 1999 and 2000.

IX.

Dooling R.J., B. Lohr, and M. L. Dent. 2000. Hearing in birds and reptiles. *In* Dooling R.J., R. R. Fay, and A. N. Popper (eds) Comparative Hearing: Birds and Reptiles. Springer-Verlag, New York, NY. Pp. 308-359.

Dudley, J. L., and K. Lajtha. 1993. The effects of prescribed burning on nutrient availability and primary production in sandplain grasslands. American Midland Naturalist 130:286-298.

Engle, D. M., R. L. Mitchell, and R. L. Stevens. 1998. Late growing-season fire effects in mid-successional tallgrass prairies. J Range Manage 51:115-121.

ESRI. 1992-1999. ArcView GIS 3.2.

Fay R. R. 1988. Hearing in vertebrates: a psychophysics databook. Hill-Fay Associates. Winnetka, IL.

Florida Fish and Wildlife Conservation Commission. 2001. Florida's Endangered species, threatened species and species of special concern. http://fcn.state.fl.us/gfc/pubs/endanger.html. Version 01FEB01.

Georgia non-game wildlife and heritage section. Georgia Natural Heritage Program. http://www.DNR.State.Ga.US/dnr/wild/. Version 01FEB01.

Gill, D. E. 1978. The metapopulation ecology of the Red-spotted Newt, *Notopthalmus viridescens* (Rafinesque). Ecological Monographs 48:145-166.

Herkert, J. R. 1994a. Breeding bird communities of midwestern prairie fragments - the effects of prescribed burning and habitat-Area. Natural Areas Journal 14:128-135.

Herkert, J. R. 1994b. The effects of habitat fragmentation on midwestern grassland bird communities. Ecological Applications 4:461-471.

Herkert, J. R. 1998. The influence of the CRP on grasshopper sparrow population trends in the mid-continental United States. Wildlife Society Bulletin 26:227-231.

Holmes, D. W. 1996. Grasshopper Sparrow (*Ammodramus savannarum*). Pages 398-399 *in* C. S. Robbins and E. A. T. Bloom, editors. Atlas of Breeding Birds of Maryland and the District of Columbia. University of Pittsburgh Press, Pittsburgh, PA.

Howe, H. F. 1994. Managing species diversity in tallgrass prairie: Assumptions and implications. Conservation Biology 8:691-704.

Hughes, J. P., R. J. Robel, K. E. Kemp, and J. L. Zimmerman. 1999. Effects of habitat on Dickcissel abundance and nest success in Conservation Reserve Program fields in Kansas. Journal of Wildlife Management 63:523-529.

Hurley, R. J., and E. C. Franks. 1976. Changes in breeding ranges of two grassland birds. Auk 93:108-115.

Iliff, M. J., R. F. Ringler, and J. L. Stasz. 1996. Field List of the Birds of Maryland, Third edition. Maryland Ornithological Society, Baltimore, MD.

Johnson, R. G., and S. A. Temple. 1986a. Assessing habitat quality for birds nesting in fragmented tallgrass prair(a)4(t)20im-11(.(l)-g-1(ve)-5(s)2456(n)]TJ 0 Tc 023-g-1962 0 Td (-)Tj 0.33

LeGrand, H. 2001. Personal communication. *In* NC Natural Heritage Program.
Loucks, O. L., M. L. Plumb-mentjes, and D. Rodgers. 1985. Gap processes and large-scale disturbances in sand prairies. *In* S. T. A. Pickett and P. S. White, editors. The ecology of natural disturbance and patch dynamics. Academic Press, San Diego, CA.
Madden, E. M., A. J. Hansen, and R. K. Murphy. 1999. Influence of prescribed fire history on habitat and abundance of passerine birds in northern mixed-grass prairie. Canadian Field-Naturalist 113:627-640.

Pennsylvania Dept. of Cons. and Nat. Res. 2001. Biota of concern in Pennsylvania. http://www.dcnr.state.pa.us/forestry/pndi/pndiweb/htm.

Peterjohn, B. G., and J. R. Sauer. 1999. Population status of North American grassland birds from the North American breeding bird survey, 1966-1996. Studies in Avian Biology 19:27-44.

Table 2a. Summary of CRP and CREP enlistment in CP2 (establishment of permanent native grasses) in Maryland Eastern Shore counties by acreage and as a percentage of total CRP and CREP enrollment 1986

Table 3. Eight species of native perennial warm-season grasses and one species of cool-season grass planted at CRFRC in March 1999.

Common Name	Scientific Name	Growing Season
Big Bluestem	Andropogon gerardi	Warm
Side-oats Grama	Bouteloua curtipendula	Warm
Deertongue Grass	Panicum clandestinum	Warm
Switch Grass D		

Table 9. Status of males of nests started in the first week of nesting season. 50% of these males were banded in 1999, but made up only 20% of 2000 male population.

Nest	Est. Lay Date	Male/Sire	Status
06-GRSP-01	5/13/00	ORBX	Banded 1999
03-GRSP-03	5/14/00	BTBX	Banded 1999
07-GRSP-02	5/14/00	YFBX	New
07-GRSP-04	5/15/00	MWBX	New
07-GRSP-01	5/16/00	YPBX	New
07-GRSP-03	5/19/00	OBBX	Banded 1999

Table 10. Nest observations from the Summer 2000 field season.

Nest	Begin Date	End Date	Days Observed	Outcome	Eggs Laid	Hatch	Fledge
2-GRSP-01	06/14/00	06/21/00	6	Fledge		5	5
2-GRSP-02	07/02/00	07/21/00	18	Fledge	4	4	4
2-GRSP-03	06/30/00	07/21/00	20	Fledge	5	4	4
2-GRSP-04	07/02/00	07/17/00	14	Fledge	5	5	5
3-GRSP-01	06/01/00	06/12/00	10	Predation	5		
3-GRSP-02	06/02/00	06/12/00	9	Fledge	4	4	4
3-GRSP-03	06/03/00	06/03/00	0	Fledge			3
3-GRSP-04	06/05/00	06/21/00	15	Fledge	5	4	3
4-GRSP-01	06/04/00	06/26/00	21	Fledge	5	5	5
4-GRSP-02	06/07/00	06/28/00	20	Fledge	5	5	5
4-GRSP-03	06/08/00	06/23/00	14	Fledge	5	5	5
4-GRSP-04	06/13/00	06/23/00	9	Predation	4		
5-GRSP-01	06/08/00	06/19/00	10	Predation	5		
5-GRSP-02	06/19/00	06/24/00	4	Fledge			5
5-GRSP-03	07/12/00	07/17/00	4	Fledge			5
6-GRSP-01	05/25/00	05/29/00	3	Fledge	5	3	
6-GRSP-02	06/21/00	07/04/00	12	Kill -mower	5	5	
6-GRSP-03	06/22/00	07/06/00	13	Fledge	4	4	4
6-GRSP-04	06/23/00	06/28/00	4	Fledge		5	5
6-GRSP-05							

Table 11. Three aspects of reproductive success: probability of nest success, number fledged per successful nest, and fecundity for 1999, 2000, and pooled data.



Figure 1. Beginning dates of nests on CRFRC as a percentage of season total based on data from 1999 (12 nests) and 2000 (30 nests). One outlier nest from late in the 1999 breeding season was omitted.

XII. APPENDIX I

We assume that: 1) all adult females (SY and ASY) are breeding each summer, 2) each female has a maximum of four opportunities to complete a nest but can physiologically complete only two nests, 3) if a nest fails a female will renest, and continue to do so until all four nest opportunities expire, and 4) the probability of daily nest survival is constant in a season.

We model the pattern of nestings through a breeding season by females as a truncated binomial probability density function of successes (p) and failures (q = 1-p) with the constraint that only two successful nests are possible:

$(p + q)^4 = p^4 + 4p^3q + 6p^2q^2 + 4pq^3 + q^4$
where the first two terms of the expansion are not possible and deleted. The remaining three terms can be described as follows;

A. Two successful nests in a season - $6p^2q^2$

1. First two nests successful, females quits further attempts = pp--
2. First nest unsuccessful, nests #2 and #3 successful, no further attempt = qpp-
3. First two nesting attempts unsuccessful, nests #3 and #4 successful = qqpp
4. First nest successful, nest # 2 not, nest # 3 successful, quit = pqp-
5. First nest successful, nests # 2 and #3 not, nest #4 successful = pqqp
6. First nest not successful, nest #2 successful, nest #3 not, nest # 4 successful = qpqp

B. One successful nest in a season - $4pq^3$

1. First nest successful, all subsequent ones failures = pqqq
2. First nest unsuccessful, second nest successful, nest #3 and #4 failures = qpqq
3. First and second nest unsuccessful, third successful, nest #4 failure = qqpq
4. First three nests unsuccessful, fourth attempt successful = qqqp

C. No successful nest in a season - q^4 = qqqq

These patterns are mutually exclusive.

D. Reproductive Success of Grasshopper Sparrows on the restored grasslands

The total number of female fledglings produced from all successful nests by a female in a season is our definition of fecundity (F). If a female has two successful nests her reproductive success is 2 x females fledglings per nest x probability of having two successful nests. A female with only one successful nest (out of 4 attempts) has reproductive success of 1 x female fledglings per nest x probability of having one successful nest. In principle, these reproductive successes are then averaged over the number of females having two, one, or zero successes.

Year	F	p	q	$6p^2q^2$	$+ 4pq^3$	$+ q^4$
1999	1.9	0.228	0.772	0.1859	0.4196	0.3552
2000	2.2	0.668	0.332	0.2951	0.0978	0.0121
Pooled	2.15	0.563	0.437	0.3632	0.1879	0.0365